

Short Note

Thamíris Christina Karlovic de Abreu*, Clarissa Alves da Rosa, Izar Aximoff and Marcelo Passamani

New record of feeding behavior by the porcupine *Coendou spinosus* (F. Cuvier, 1823) in high-altitude grassland of the Brazilian Atlantic Forest

DOI 10.1515/mammalia-2016-0026

Received February 19, 2016; accepted July 6, 2016

Abstract: *Coendou spinosus* is a rodent from the Brazilian Atlantic Forest with a cryptic behavior that hinders collection of information regarding its ecology. On October 2014, we observed an individual of *C. spinosus* above 2100 m of altitude, on the high-altitude grasslands of the Itatiaia National Park, feeding on the flowers of *Camptosema scarlatinum*. This is the highest altitude ever recorded and the first registration of vine species consumption and ground feeding behavior by the *C. spinosus*. Therefore, we recommend the undertaking of new studies to understand its ecological requirements and interspecific interactions.

Keywords: diet; Erethizontidae; florivory; herbivory; Itatiaia National Park.

The Erethizontidae family forms two ecological groups of New World porcupines, the North American porcupine *Erethizon dorsatum* (C. Linnaeus, 1758), who's weight exceeds 10 kg and is both tree and ground-dwelling (Woods 1973, Roze 2012, Voss 2015); and the South American porcupines (*Chaetomys subspinosus* I. Olfers, 1818 and *Coendou* sp.), that weigh up to 5 kg, and are known

for their prehensile-tail, conducting their daily activities mainly in trees (Chiarello et al. 1997, Giné et al. 2010, Souto-Lima et al. 2010, Voss 2015). All the species of Erethizontidae feed mainly on leaves, fruit and bark (Roze 2012). The studies of South American porcupines focus on *C. subspinosus*, an endangered species with high ecological requirements (Chiarello et al. 1997, Giné et al. 2010, 2012, 2015, Souto-Lima et al. 2010, Zortéa and Brito 2010, Oliveira et al. 2012), but little is known about the *Coendou* species because of their cryptic habits that hinder acquisition of knowledge about their ecological aspects.

The porcupine *Coendou spinosus* (F. Cuvier, 1823) is a nocturnal species which occupies areas up to 900 m of altitude in subtropical forests (Voss 2011). In a recovering forest fragment surrounded by anthropogenic habitats, Passamani (2010) studied a single individual that showed a diet basically folivore and feeding in trees. In this paper we provide new information about the *C. spinosus* feeding behavior and habitat based on a single observation in the Itatiaia National Park (INP), Brazil.

Located in the states of Minas Gerais and Rio de Janeiro, the INP has 28,084 ha of Brazilian Atlantic Forest (Barreto et al. 2013) and is considered irreplaceable for biodiversity conservation (Le Saout et al. 2013). The INP has three vegetation types: montane forest (from 500 to 1500 m); upper montane forest (from 1500 to 2000 m); and high-altitude grasslands (from 1800 to 2791 m) (Segadas-Vianna and Dau 1965). With a mesothermic climate (Köppen 1936), mean annual temperature and precipitation of around 11.5°C and 2150 m, respectively (Barreto et al. 2013), the high-altitude grasslands dominate the landscape above 2000 m and have a predominance of herbaceous plants, shrubs and small trees that can reach 4 m high (Martinelli 1996, Aximoff 2011). On October 8th, 2014, at 2137 m, on high-altitude grasslands of INP (22°24'12"S and 44°38'34"W, WGS 84) (Figure 1), around 8:00 am, we observed an individual of *Coendou*

*Corresponding author: Thamíris Christina Karlovic de Abreu, Laboratório de Ecologia e Conservação de Mamíferos, Setor de Ecologia, Departamento de Biologia, Universidade Federal de Lavras, Campus Universitário, Lavras, Minas Gerais, 37200-000, Brazil, e-mail: thamycka@hotmail.com

Clarissa Alves da Rosa and Marcelo Passamani: Laboratório de Ecologia e Conservação de Mamíferos, Setor de Ecologia, Departamento de Biologia, Universidade Federal de Lavras, Campus Universitário, Lavras, Minas Gerais, 37200-000, Brazil

Izar Aximoff: Instituto de Pesquisas do Jardim Botânico do Rio de Janeiro, Escola Nacional de Botânica Tropical, Pós-Graduação em Botânica. Rua Pacheco Leão 2040 – Solar da Imperatriz, Horto – Rio de Janeiro, RJ, 22460-036, Brazil




Figure 1: Study area showing the Itatiaia National Park area and the registration point of *Coendou spinosus* feeding on *Camptosema scarlatinum*.

spinosus feeding on flowers of *Camptosema scarlatinum* (Mart. ex Benth.) Burkart (Fabaceae) (Figure 2C and D). The porcupine was on the ground using both forefeet to bring the flowers into its mouth. After our presence was noticed, it started an escape behavior and climbed a shrub (approx. 2 m high) next to the feeding site (Figure 2A and B).

The South American porcupines are known to feed in trees (Giné et al. 2010, Souto-Lima et al. 2010) and our record demonstrates a new feeding behavior in these species. However, the *Coendou* species, including *C. spinosus*, are frequently killed on Brazilian roads (e.g. Freitas et al. 2013), showing that they descend to the ground more than initially expected. Usually, this ground exposure is most observed when the porcupine descends to the ground to defecate or urinate (Chiarello et al. 1997, Passamani 2010, Oliveira et al. 2012) and even then, it is undertaken with caution because of the greater exposure to predation (Oliveira et al. 2012).

Also, our record is the first consumption of vine species by *Coendou spinosus*, since only six species of trees are known to be feed on by the species (Passamani 2010). *Camptosema scarlatinum* is a terrestrial liana that

can be found on grasslands and at forest edges of Brazilian savannah and Atlantic forest (Burkart 1970). Considering that our registration was made in an open grassland, *C. scarlatinum* can be an alternative food item that is not important in forested habitats where other tree products are available (Roze 2012).

Coendou spinosus occurs in forest fragments of different conservation levels. Although, previous registrations in INP showed only *C. insidiosus* (I. Olfers, 1818) in Montane forest of up to 1200 m (Ávila-Pires and Gouvêa 1977, Geise et al. 2004). Our registration in an open grassland habitat above 2100 m of altitude is the highest altitude of occurrence at which *C. spinosus* has ever been recorded. The high altitude grasslands concentrate most of the endemic species and receives the lowest human pressure of the INP (Geise et al. 2004, Cunha 2010, Barreto et al. 2013), including recent records of primates and carnivore species (Rosa et al. 2015, Aximoff and Vaz 2016). We do not know the habitat use and behavior of the porcupines in those environments, but some authors suggest that food resources in high-altitude environments may be more abundant than previously thought (Raboy et al. 2013, Aximoff 2015). Perhaps *C. spinosus*




Figure 2: *Coendou spinosus* on a shrub next to the feeding site (A and B); *Camptosema scarlatinum* flourishing in the high altitude grassland of the Itatiaia National Park (C); and *Camptosema scarlatinum* flower (D).

use the forest patches that are immersed in the grassland matrix or, because of high altitude, even change the food items preferences during winter, as occurs with *Erythizon dorsatum* at high latitudes (Roze 2012). To answer these questions we need the development of systematic studies regarding habitat use, diet and food resources availability for *C. spinosus*. Considering that other species of the South American porcupines (e.g. *C. subspinosus*) are highly selective in terms of species consumed (e.g. Giné et al. 2010) we also need studies about the plant-animal interaction in order to understand ecological needs and interspecific interactions of the *Coendou* species.

Acknowledgments: We would like to thank Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG) and Tropical Forest Conservation Act (TFCA/FUNBIO) for financial resources. We thank the Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio – license 36838), and the administration of the Itatiaia National Park for logistical support during this study. We also thank Mr. José Cristiano for supporting our expeditions.

References

- Ávila-Pires, F.D. and E. Gouvêa. 1977. Mamíferos do Parque Nacional do Itatiaia. Boletim do Museu Nacional, Rio de Janeiro 291: 1–29.
- Aximoff, I. 2011. O que perdemos com a passagem do fogo pelos campos de altitude do Estado do Rio de Janeiro. Revista Biodiversidade Brasileira, Número Temático: Ecologia e Manejo do Fogo em Áreas Protegidas. 2: 180–200.
- Aximoff, I. 2015. Confirmação da ocorrência do muriqui-do-norte (Primates, Atelidae) no Parque Nacional do Itatiaia, Estado do Rio de Janeiro, sudeste do Brasil. Oecologia Australis 18: 1–5.
- Aximoff, I. and S.M. Vaz. 2016. Bugio-ruivo (Primates, Atelidae) em campos de altitude e com anomalia na coloração no Parque Nacional do Itatiaia, sudeste do Brasil. Oecologia Australis 20: 122–127.
- Barreto, C.G., J.B. Campos, D.M. Roberto, D.M. Roberto, N.T. Schwarze, G.S.G. Alves and W. Coelho. 2013. Análise da Região da Unidade de Conservação. In: Plano de Manejo Parque Nacional do Itatiaia, Encartes 2 e 3. Instituto Chico Mendes de Conservação da Biodiversidade, Brasília. pp. 1–117.
- Burkart, A. 1970. Las leguminosas Faseóleas argentinas de los géneros *Mucuna*, *Dioclea* y *Camptosema*. Darwiniana 16: 175–217.
- Chiarello, A.G., M. Passamani and M. Zortéa. 1997. Field observations on the thin-spined porcupine, *Chaetomys subspinosus* (Rodentia: Echimyidae). Mammalia 61: 29–36.
- Cunha, A.A. 2010. Negative effects of tourism in a Brazilian Atlantic forest National Park. J. Nat. Conserv. 4: 291–295.

- Freitas, M.A., D.P.F. França and D. Veríssimo. 2013. First record of the Bicoloured-spined Porcupine *Coendou bicolor* (Tschudi, 1884) for Brazil. Check List 9: 94–96.
- Geise, L., L.G. Pereira, D.E.P. Bossi and H.G. Bergallo. 2004. Pattern of elevational distribution and richness of non-volant mammals in Itatiaia National Park and its surroundings, in Southeastern Brazil. Braz. J. Biol. 64: 599–612.
- Giné, G.A.F., J.M.B. Duarte and D. Faria. 2010. Feeding ecology of a selective folivore, the thin-spined porcupine (*Chaetomys subspinosus*) in the Atlantic forest. J. Mammal. 91: 931–941.
- Giné, G.A.F., J.M.B. Duarte, T.C.S. Motta and D. Faria. 2012. Activity, movement and secretive behavior of threatened arboreal folivore, the thin-spined porcupine, in the Atlantic forest of southern Bahia, Brazil. J. Zool. 286: 131–139.
- Giné, G.A.F., E.H. Barros, J.M.B. Duarte and D. Faria. 2015. Home range and multiscale habitat selection of threatened thin-spined porcupine in the Brazilian Atlantic Forest. J. Mammal. 96: 1095–1105.
- Köppen, W. 1936. Das geographische System der Klimate. In: (W. Koppen and R. Geiger, eds.) Klimatologie. Gebr, Borntraeger, Germany. pp. 1–44.
- Le Saout, S., M. Hoffmann, Y. Shi, A. Hughes, C. Bernard, T.M. Brooks, B. Bertzky, S.H.M. Butchart, S.N. Stuart, T. Badman and A.S.L. Rodrigues. 2013. Protected areas and effective biodiversity conservation. Science 342: 803–805.
- Martinelli, G. 1996. Campos de Altitude. Index, Rio de Janeiro. pp. 160.
- Oliveira, P.A., R.B. Souto-Lima and A.G. Chiarello. 2012. Home range, movements and diurnal roosts of the endangered thin-spined porcupine, *Chaetomys subspinosus* (Rodentia: Erethizontidae), in the Brazilian Atlantic Forest. Mammal. Biol. 77: 97–107.
- Passamani, M. 2010. Use of space and activity pattern of *Sphiggurus villosus* (F. Cuvier, 1823) from Brazil (Rodentia: Erethizontidae). Mammal. Biol. 75: 455–458.
- Raboy, B.E., L.G. Neves, S.L. Zeigler and L.C. Oliveira. 2013. Occurrences of the Golden-headed Lion Tamarin (*Leontopithecus chrysomelas*) above 500 meters in Southern Bahia, Brazil and implications for conservation planning. Primate Conserv. 1: 25–31.
- Rosa, C.A.R., K.K. Santos, G.M.M. Faria, F. Puertas and M. Pasamani. 2015. Diet of *Chrysocyon brachyurus* (Illiger, 1815) in Southeastern Brazil. Bol. Soc. Bras. Mastozool. 72: 7–10.
- Roze, U. 2012. Porcupines: the animal answer guide. John Hopkins University Press, Baltimore.
- Segadas-Vianna, F. and L. Dau. 1965. Ecology of Itatiaia range, southeastern Brazil: II Climates and altitudinal climatic zonation. Arquivos do Museu Nacional 53: 31–53.
- Souto-Lima, R.B., P.A. Oliveira and A.G. Chiarello. 2010. Diet of the thin-spined porcupine (*Chaetomys subspinosus*), an Atlantic forest endemic threatened with extinction in southeastern Brazil. Mammal. Biol. 75: 538–546.
- Voss, R.S. 2011. Revisionary notes on Neotropical porcupines (Rodentia: Erethizontidae) 3. An annotated checklist of the species of *Coendou* Lacépède, 1799. American Museum Novitates 3720: 1–36.
- Voss, R.S. 2015. Superfamily Erethizontoidea Bonaparte, 1845. In: (J.L. Patton, U.F.J. Pardiñas and G. D'Elía, eds.) Mammals of South America, Volume 2: Rodents. The University of Chicago Press, Chicago and London. pp. 786–805.
- Woods, C.A. 1973. *Erethizon dorsatum*. Mamm. Species 29: 1–6.
- Zortéa, M. and B.F.A. de Brito. 2010. Diurnal roosts and minimum home range defined by sleeping sites of a thin-spined porcupine *Chaetomys subspinosus* (Rodentia: Erethizontidae). Zootaxa 27: 209–212.